Insomnia and Naturopathy: An Evidence-Based Review of Non-Pharmacological Approaches to Sleep Management

Priyanka Gupta
Department of Psychology
Renaissance University Indore, Indore (M.P.), India
https://doi.org/10.64882/ijrt.v14.iS2.2050

Abstract
Insomnia is a prevalent sleep disorder characterised by persistent difficulty initiating sleep, maintaining sleep, or gaining restorative sleep together with meaningful daytime impairment. Interest in non-pharmacological and complementary approaches is increasing because chronic insomnia often requires long-term management and because many patients prefer strategies that can be incorporated into daily life. This narrative review evaluates naturopathic and complementary approaches to insomnia that could be within the current evidence-based treatment framework. The strongest evidence supports multicomponent cognitive behavioural therapy for insomnia (CBT-I), which is recommended as first-line treatment in major sleep-medicine guidelines. Adjunctive approaches with varying le0vels of support include regular exercise, yoga and other mind-body practices, mindfulness, appropriately timed warm bathing, circadian-light management, aromatherapy, and acupuncture. Evidence for magnesium, valerian, chamomile, and other natural products is inconsistent or limited, while detoxification procedures and colon hydrotherapy lack convincing evidence for insomnia treatment. Overall, naturopathic care is most defensible when it emphasises low-risk lifestyle and behavioural measures, individualised assessment, and integration with established insomnia care rather than replacement of evidence-based treatment. Future studies should use standardised insomnia diagnoses, validated outcomes, adequate sample sizes, active controls, objective sleep measures where appropriate, and longer follow-up.

Keywords: Insomnia; naturopathy; non-pharmacological treatment; CBT-I; yoga; mindfulness; exercise; hydrotherapy; aromatherapy; acupuncture; sleep hygiene

Introduction

Insomnia is one of the most frequently encountered sleep disorders. It is clinically important because impaired sleep can coexist with reduced daytime functioning, fatigue, mood symptoms, cognitive complaints, and lower quality of life. A 2025 meta-analysis of population studies that used diagnostic approaches consistent with DSM criteria estimated that insomnia disorder affects a substantial minority of adults, while prevalence varies according to the method of assessment and population studied. The disorder is now understood as more than simply "too little sleep": persistent cognitive-emotional and physiological arousal, maladaptive sleep-related behaviours, circadian factors, comorbid illness, and environmental influences may all contribute to its onset or maintenance. [1][2]

Naturopathy traditionally emphasises lifestyle, behavioural, nutritional, physical, and mind-body approaches intended to support self-regulation and general health. In sleep care, this orientation overlaps with several components of modern behavioural sleep medicine, such as regular routines, physical activity, stress reduction, and attention to environmental cues. However, a scientifically responsible review must distinguish between interventions with strong evidence for chronic insomnia and those for which evidence may be preliminary, inconsistent, or absent.

Current clinical guidelines provide an essential reference point. The American Academy of Sleep Medicine (AASM) strongly recommends multicomponent CBT-I for adults with chronic insomnia; relaxation therapy receives a conditional recommendation and sleep hygiene alone is not recommended as a stand-alone treatment. The 2023 European insomnia guideline similarly identifies CBT-I as the first-line treatment for chronic insomnia in adults of all ages, including those with comorbidities. Therefore, complementary and naturopathic interventions are best evaluated as potential adjuncts to, or supportive elements within, comprehensive insomnia care rather than as substitutes for established treatment. [3][4]

Aim and Review Method

This review aims to examine commonly used naturopathic and complementary interventions for insomnia, evaluate the strength and overall direction of the available evidence, explore their possible mechanisms of action, highlight relevant safety considerations and evidence gaps, and clarify how these approaches can be integrated with guideline-based management.

This article is a narrative synthesis of selected clinical practice guidelines, systematic reviews, meta-analyses, and relevant clinical studies on insomnia and complementary sleep interventions. It considers behavioural treatment, exercise, yoga and Yoga Nidra, mindfulness, passive body heating, light exposure, aromatherapy, acupuncture, magnesium, and herbal products. The selection is not presented as a reproducible systematic search: no registered protocol, comprehensive screening record, or formal independent risk-of-bias assessment is reported. Evidence descriptions reflect the findings and limitations of the cited sources rather than a new pooled analysis. Conclusions are therefore interpretive and should not be taken as an exhaustive assessment of all available studies.

Evidence-Based Framework for Non-Pharmacological Management

CBT-I, Behavioural Treatment, and Sleep Hygiene

CBT-I combines several components—commonly stimulus control, sleep restriction or sleep-compression strategies, cognitive restructuring, relaxation, and sleep-related education—to target the behavioural and cognitive processes that perpetuate insomnia. Guideline recommendations are stronger for multicomponent CBT-I than for any single lifestyle measure. This distinction matters because "sleep hygiene" is often used broadly in wellness and naturopathic settings; sleep hygiene education by itself has not shown sufficient efficacy to be recommended as a stand-alone treatment for chronic insomnia. [3][4]

In a naturopathic model, sleep hygiene can still serve as a supportive foundation. Useful measures include maintaining a consistent sleep-wake schedule, limiting caffeine later in the day, avoiding heavy meals and excessive alcohol close to bedtime, minimising disruptive light

and device use before sleep, creating a dark and quiet sleep environment, and separating the bed from prolonged wakeful activities. These measures are most appropriate when incorporated into a broader behavioural plan rather than presented as a complete treatment for persistent insomnia.

Relaxation and the Hyperarousal Model

Hyperarousal is a central concept in contemporary models of insomnia. It includes cognitive-emotional activation such as worry and rumination as well as cortical and, in some studies, physiological arousal. This supports the use of relaxation-based strategies aimed at lowering physiological and mental arousal before sleep; however, a theoretically sound mechanism alone is not sufficient to confirm clinical effectiveness. [2]

Relaxation practices may include progressive muscle relaxation, controlled breathing, guided imagery, and other structured techniques. The AASM guideline gives relaxation therapy a conditional recommendation as a single-component treatment, which supports its use when CBT-I is unavailable, unacceptable, or being delivered in modular form. Massage and similar body-based practices may improve subjective relaxation, but the evidence base for treating chronic insomnia specifically is smaller than that for CBT-I. [3]

Physical Activity and Exercise

Regular exercise is one of the more defensible lifestyle interventions for sleep. A systematic review focused on adults with insomnia found improvement in subjective sleep quality, although methodological limitations reduced confidence in several outcomes. A later meta-analysis of randomized controlled trials also found that regular physical and mind-body exercise improved subjective sleep outcomes, with more limited changes in objective sleep measures. More recent evidence continues to support a modest beneficial role for exercise, while also showing substantial heterogeneity across intervention types, intensity, duration, and participant characteristics. [5][6]

From a practical perspective, exercise may support sleep through several pathways: regulation of arousal, mood improvement, increased homeostatic sleep pressure, cardiometabolic benefits, and reinforcement of regular daily rhythms. Moderate aerobic activity and resistance exercise can be individualized according to age, fitness, comorbidities, and patient preference. Excessively vigorous activity immediately before bedtime may be stimulating for some individuals, so timing should be individualized.

Yoga, Pranayama, Yoga Nidra, and Mindfulness

Yoga and related mind-body practices combine physical movement, breath regulation, attentional control, and relaxation. These characteristics make them particularly relevant to insomnia characterized by stress, rumination, or somatic tension. Systematic reviews suggest that mind-body exercise can improve subjective sleep quality, but intervention protocols vary widely and many studies are small or at risk of bias. [6][7]

Mindfulness-based interventions have shown improvement in some insomnia and sleep-quality outcomes compared with nonspecific controls, although the size and interpretation of benefits depend on the outcome and comparator. Therefore, mindfulness is best understood as an

adjunct that may reduce pre-sleep rumination and support attentional regulation rather than as a replacement for CBT-I. [8]

Yoga Nidra has received growing research attention. A 2026 systematic review and meta-analysis found potentially meaningful improvements in sleep-related outcomes, but the pooled randomized evidence was small, heterogeneous, and rated as very low certainty. The authors emphasized that current findings should be considered preliminary. This is especially relevant in naturopathy and yoga-based practice: Yoga Nidra can reasonably be offered as a low-risk relaxation adjunct while avoiding claims of proven efficacy for chronic insomnia. [7]

Hydrotherapy and Pre-Sleep Passive Body Heating

Warm-water bathing is one of the clearest examples in which a traditional or naturopathic practice has a plausible physiological mechanism and supportive quantitative evidence. A systematic review and meta-analysis of water-based passive body heating reported that a warm bath or shower in the range of approximately 40–42.5°C, when timed about one to two hours before bedtime, was associated with improved self-rated sleep quality and sleep efficiency and could shorten sleep-onset latency. The proposed mechanism is enhanced peripheral heat loss after bathing, which facilitates the normal decline in core body temperature associated with sleep initiation. [9]

This evidence does not justify broad claims that "hydrotherapy" cures insomnia; rather, it supports a specific, timed thermal intervention. Individuals with cardiovascular disease, autonomic instability, pregnancy-related concerns, mobility limitations, or risk of falls should use heat exposure cautiously and seek appropriate clinical advice.

Light Exposure and Circadian Regulation

Light is the dominant environmental signal for entraining human circadian rhythms. Consistent wake times and adequate daytime, especially morning, light exposure can help stabilize sleep-wake timing, while excessive bright or blue-enriched light late in the evening may delay sleep in susceptible individuals. Light-based interventions are particularly relevant when insomnia symptoms coexist with delayed sleep timing or other circadian rhythm disturbance. The 2023 umbrella review of complementary and alternative treatments found that the evidence for light exposure in insomnia disorder was less convincing than for some other modalities, illustrating the importance of matching light interventions to the patient's circadian phenotype rather than applying them indiscriminately. [10]

Aromatherapy

Aromatherapy, particularly lavender-based inhalation, is widely used for relaxation. Meta-analyses have reported improvements in subjective sleep quality, but the literature is characterized by substantial heterogeneity, variable product composition and delivery methods, small studies, and risk of publication bias. These limitations mean that aromatherapy should be described as a potentially useful comfort or relaxation adjunct rather than a well-established insomnia treatment. [11]

Safety is generally favourable when essential oils are used appropriately, but concentrated oils may cause skin irritation, allergic reactions, headache, or respiratory discomfort in sensitive

individuals. Oral ingestion of essential oils should not be assumed safe simply because a product is described as "natural."

Acupuncture

Acupuncture has been evaluated in numerous trials and systematic reviews. Meta-analyses have reported improvements in subjective measures such as the Pittsburgh Sleep Quality Index and, in some analyses, selected objective sleep measures. However, many trials have substantial risk of bias, high heterogeneity, variable comparators, and challenges with blinding. Consequently, acupuncture may be considered an adjunctive option for some patients, but the certainty of evidence remains lower than for CBT-I. [12][10]

Diet, Magnesium, and Herbal Products

Nutrition influences general health and may affect sleep through metabolic, circadian, and neurotransmitter pathways, but evidence for specific nutrients as treatments for chronic insomnia is limited. A balanced diet, regular meal timing, avoidance of excessive caffeine, and avoidance of very heavy late-night meals are reasonable components of sleep-supportive care. Claims that isolated nutrients reliably treat insomnia require stronger evidence.

Magnesium is commonly marketed for sleep. A 2021 systematic review in older adults found a reduction in sleep-onset latency in pooled data, but only three small randomized trials were available, all with moderate-to-high risk of bias, and the overall certainty of evidence was low to very low. Magnesium therefore should not be represented as a routine evidence-based treatment for insomnia, although deficiency correction may be clinically appropriate when indicated. [13]

Herbal products also require caution. Earlier systematic review evidence did not establish a clear benefit of herbal medicines such as valerian and chamomile over placebo for insomnia, largely because trials were small and methodologically inconsistent. A 2024 umbrella review of valerian concluded that evidence for treating insomnia remained weak or inconclusive, although some subjective sleep-quality outcomes may improve. Chamomile may improve some sleep-quality outcomes in pooled analyses, but direct evidence for insomnia severity remains limited. Product standardization, dose, interactions, contamination, and adverse effects are additional concerns. [14][15][16]

Detoxification, Fasting, Mud Therapy, Colon Hydrotherapy, and Shirodhara

Some naturopathic systems describe fasting, mud applications, or colon hydrotherapy as "detoxification." These practices should not be presented as established insomnia treatments on the basis of the evidence reviewed here. The National Center for Complementary and Integrative Health notes that research on detoxification programmes is limited and often methodologically weak. Restrictive diets may provide inadequate nutrition; fasting can cause weakness and dehydration; and colon-cleansing procedures can have serious adverse effects. Such general safety information does not establish efficacy for insomnia. [17]

Shirodhara is a traditional Ayurvedic oil-dripping procedure. A small randomized crossover pilot study compared sesame-oil Shirodhara with warm-water Shirodhara in adults reporting sleep problems. Its size and comparator limit conclusions about treatment of diagnosed chronic

insomnia and about effects beyond warmth, attention, and the procedure itself. The study provides preliminary evidence rather than proof of established efficacy. [18]

Comparative Summary of Evidence

Intervention	Current evidence	Reasonable role	Key caution
CBT-I	Strong guideline support	First-line treatment for chronic insomnia	Requires trained delivery or validated digital program
Sleep hygiene alone	Insufficient as stand-alone treatment	Supportive foundation within broader treatment	Do not present as complete treatment
Relaxation therapy	Conditional guideline recommendation	Adjunct or modular behavioural treatment	Effect smaller/less certain than multicomponent CBT-I
Exercise	Moderate evidence for subjective sleep improvement	Routine lifestyle adjunct	Individualize intensity and timing
Yoga/mind-body practice	Promising; heterogeneous evidence	Adjunct for stress, arousal, flexibility, wellbeing	Protocols and study quality vary
Mindfulness	Promising adjunct	Reduce rumination and stress reactivity	Outcomes and certainty depend on comparator and study quality
Yoga Nidra	Preliminary, very low-certainty evidence	Relaxation adjunct	Small, geographically concentrated evidence base
Warm bath/shower	Supportive meta-analytic evidence for timed passive heating	Simple pre-sleep adjunct	Heat/fall/cardiovascular precautions
Light management	Mechanistically important; insomnia evidence context-dependent	Circadian stabilization when timing is appropriate	Incorrect timing can worsen phase delay/advance

Intervention	Current evidence	Reasonable role	Key caution
Aromatherapy	Possible subjective benefit; high heterogeneity	Comfort/relaxation adjunct	Allergy, irritation, variable formulations
Acupuncture	Possible benefit; risk of bias and heterogeneity	Optional adjunct	Quality of evidence below CBT-I
Magnesium	Low/very low-certainty evidence	Correct deficiency; not routine insomnia therapy	Interactions, renal impairment, GI effects
Valerian/chamomile	Mixed or limited evidence	Optional carefully selected adjunct	Standardization, interactions, inconsistent efficacy
Detox/colon hydrotherapy	No established insomnia efficacy	Not recommended as insomnia treatment	Potential direct harms and unsupported mechanism

Clinical Integration and Safety Considerations

A practical evidence-based naturopathic approach begins with assessment rather than immediate selection of a remedy. Persistent insomnia may coexist with sleep apnoea, restless legs syndrome, circadian rhythm disorders, depression, anxiety, chronic pain, endocrine disease, medication effects, substance use, menopausal symptoms, or other medical conditions. Symptoms such as loud snoring with witnessed apnoeas, marked daytime sleepiness, unusual nocturnal movements, severe mood symptoms, or refractory insomnia warrant formal medical or sleep evaluation.

For uncomplicated chronic insomnia, a stepped integrative plan can prioritize CBT-I or its validated digital forms, while adding regular exercise, circadian regularity, relaxation or mindfulness, and an appropriately timed warm bath when acceptable to the individual. Yoga may provide both physical and psychological benefit. Aromatherapy or acupuncture may be considered for preference-sensitive adjunctive use after discussing the lower certainty of evidence. Nutritional supplements and herbs should be reviewed for dose, quality, interactions, contraindications, and realistic expectations.

The term "natural" should not be equated with "risk free." Supplement dose, product quality, medication interactions, renal function, and other individual contraindications require review. Magnesium trials provide limited safety data, and herbal evidence is affected by variable preparations and study methods. Integration with conventional care is particularly important when symptoms persist or comorbid disease complicates management. [13–17]

Limitations of the Current Evidence

The evidence base for complementary and naturopathic sleep interventions has several recurring limitations. Many studies are small, use short treatment periods, rely heavily on self-reported sleep quality, and include participants with general sleep complaints rather than rigorously diagnosed insomnia disorder. Blinding is difficult for behavioural, yoga, acupuncture, and aromatherapy interventions. Control groups vary from wait-list to placebo-like procedures or active treatment, making effect sizes difficult to compare directly. Intervention protocols are also heterogeneous; "yoga," "mindfulness," "aromatherapy," and "acupuncture" can refer to substantially different practices. Finally, long-term safety, adherence, durability of benefit, and comparative effectiveness against CBT-I are often inadequately studied. [10]

These limitations reinforce a graded interpretation: a therapy can be biologically plausible and associated with improved subjective sleep without yet having the evidence required to call it an established treatment for chronic insomnia.

Future Research Directions

Future research should use standardized diagnostic criteria for insomnia disorder, preregistered protocols, adequate sample sizes, transparent randomization, credible control conditions, validated measures such as the Insomnia Severity Index and sleep diaries, and objective measures such as actigraphy or polysomnography when clinically relevant. Studies should report adverse events and adherence, specify intervention dose and timing, and include longer follow-up. Comparative trials examining adjunctive naturopathic interventions added to CBT-I would be particularly useful because this design reflects realistic integrative clinical practice. For traditional interventions such as Yoga Nidra, Shirodhara, and hydrotherapeutic practices, research should distinguish clearly between the cultural/traditional rationale and the tested biomedical mechanism. This would allow traditional practices to be studied respectfully while avoiding unsupported claims.

Conclusion

Naturopathic and complementary approaches can contribute meaningfully to a patient-centred strategy for sleep when they are selected according to the strength of evidence and individual clinical context. The best-supported non-pharmacological treatment for chronic insomnia remains CBT-I. Regular exercise, relaxation, yoga and other mind-body practices, mindfulness, circadian-light management, and timed warm bathing can serve as practical adjuncts, although the certainty and magnitude of benefit differ across interventions. Aromatherapy and acupuncture may improve subjective sleep outcomes for some individuals but are supported by more heterogeneous evidence. Magnesium, valerian, chamomile, and other natural products should be discussed cautiously because results are inconsistent and product-related safety issues remain. Detoxification procedures, colon hydrotherapy, and other poorly studied practices should not be represented as established insomnia treatments. A modern naturopathic approach is therefore strongest when it combines lifestyle-oriented care with transparent evidence grading, safety screening, and appropriate referral or integration with guideline-based sleep medicine.

References

1. van Straten A, Weinreich KJ, Fábián B, Reesen JE, Grigori S, Luik AI, Harrer M, Lancee J. The prevalence of insomnia disorder in the general population: a meta-analysis. Journal of Sleep Research. 2025;34(5):e70089. doi:10.1111/jsr.70089.
2. Dressle RJ, Riemann D. Hyperarousal in insomnia disorder: current evidence and potential mechanisms. Journal of Sleep Research. 2023;32(6):e13928. doi:10.1111/jsr.13928.
3. Edinger JD, Arnedt JT, Bertisch SM, Carney CE, Harrington JJ, Lichstein KL, Sateia MJ, Troxel WM, Zhou ES, Kazmi U, Heald JL, Martin JL. Behavioral and psychological treatments for chronic insomnia disorder in adults: an American Academy of Sleep Medicine clinical practice guideline. Journal of Clinical Sleep Medicine. 2021;17(2):255-262. doi:10.5664/jcsm.8986.
4. Riemann D, Espie CA, Altena E, et al. The European Insomnia Guideline: an update on the diagnosis and treatment of insomnia 2023. Journal of Sleep Research. 2023;32(6):e14035. doi:10.1111/jsr.14035.
5. Lowe H, Haddock G, Mulligan LD, Gregg L, Fuzellier-Hart A, Carter LA, Kyle SD. Does exercise improve sleep for adults with insomnia? A systematic review with quality appraisal. Clinical Psychology Review. 2019;68:1-12. doi:10.1016/j.cpr.2018.11.002.
6. Xie Y, Liu S, Chen XJ, Yu HH, Yang Y, Wang W. Effects of exercise on sleep quality and insomnia in adults: a systematic review and meta-analysis of randomized controlled trials. Frontiers in Psychiatry. 2021;12:664499. doi:10.3389/fpsyt.2021.664499.
7. Singh S, Dhanlika D, Singh U, Haldar P, Sharma G. Effect of Yoga Nidra (yogic sleep) in sleep disturbances and insomnia: systematic review and metanalysis. Sleep and Breathing. 2026;30(2):144. doi:10.1007/s11325-026-03685-0.
8. Wang YY, Wang F, Zheng W, Zhang L, Ng CH, Ungvari GS, Xiang YT. Mindfulness-based interventions for insomnia: a meta-analysis of randomized controlled trials. Behavioral Sleep Medicine. 2020;18(1):1-9. doi:10.1080/15402002.2018.1518228.
9. Haghayegh S, Khoshnevis S, Smolensky MH, Diller KR, Castriotta RJ. Before-bedtime passive body heating by warm shower or bath to improve sleep: a systematic review and meta-analysis. Sleep Medicine Reviews. 2019;46:124-135. doi:10.1016/j.smrv.2019.04.008.
10. Ell J, Schmid SR, Benz F, Spille L. Complementary and alternative treatments for insomnia disorder: a systematic umbrella review. Journal of Sleep Research. 2023;32(6):e13979. doi:10.1111/jsr.13979.
11. Hwang E, Shin S. The effects of aromatherapy on sleep improvement: a systematic literature review and meta-analysis. Journal of Alternative and Complementary Medicine. 2015;21(2):61-68. doi:10.1089/acm.2014.0113.
12. Shergis JL, Ni X, Jackson ML, Zhang AL, Guo X, Li Y, Lu C, Xue CC. A systematic review of acupuncture for sleep quality in people with insomnia. Complementary Therapies in Medicine. 2016;26:11-20. doi:10.1016/j.ctim.2016.02.007.

13. Mah J, Pitre T. Oral magnesium supplementation for insomnia in older adults: a systematic review and meta-analysis. BMC Complementary Medicine and Therapies. 2021;21(1):125. doi:10.1186/s12906-021-03297-z.
14. Leach MJ, Page AT. Herbal medicine for insomnia: a systematic review and meta-analysis. Sleep Medicine Reviews. 2015;24:1-12. doi:10.1016/j.smrv.2014.12.003.
15. Valente V, Machado D, Jorge S, Drake CL, Marques DR. Does valerian work for insomnia? An umbrella review of the evidence. European Neuropsychopharmacology. 2024;82:6-28. doi:10.1016/j.euroneuro.2024.01.008.
16. Kazemi A, Shojaei-Zarghani S, Eskandarzadeh P, Hashempur MH. Effects of chamomile (Matricaria chamomilla L.) on sleep: a systematic review and meta-analysis of clinical trials. Complementary Therapies in Medicine. 2024;84:103071. doi:10.1016/j.ctim.2024.103071.
17. National Center for Complementary and Integrative Health. "Detoxes" and "cleanses": what you need to know. Updated March 2025. Accessed October 24, 2026. https://www.nccih.nih.gov/health/detoxes-and-cleanses-what-you-need-to-know.
18. Tokinobu A, Yorifuji T, Tsuda T, Doi H. Effects of Ayurvedic oil-dripping treatment with sesame oil vs. with warm water on sleep: a randomized single-blinded crossover pilot study. Journal of Alternative and Complementary Medicine. 2016;22(1):52–58. doi:10.1089/acm.2015.0018.